

Applications of double int.

Mass, Centers of Mass, and Double Integrals

(19)

Suppose a 2-D region R has density $\rho(x, y)$ at each point (x, y) . We can partition R into subrectangles, with m of them in the x -direction, and n in the y -direction. Suppose each subrectangle has width Δx and height Δy . Then a subrectangle containing the point (\hat{x}, \hat{y}) has approximate mass

$$\rho(\hat{x}, \hat{y})\Delta x\Delta y$$

and the mass of R is approximately

$$\sum_{i=1}^m \sum_{j=1}^n \rho(x_i, y_j)\Delta x\Delta y$$

where (x_i, y_j) is a point in the i, j -th subrectangle. Letting m and n go to infinity, we have

$$M = \text{mass of } R = \iint_R \rho(x, y) dA.$$

Similarly, the moment with respect to the x axis can be calculated as

$$M_x = \iint_R y\rho(x, y) dA$$

and the moment with respect to the y axis can be calculated as

$$M_y = \iint_R x\rho(x, y) dA.$$

The we may calculate the center of mass of R via

$$\text{center of mass of } R = (\bar{x}, \bar{y}) = \left(\frac{M_y}{M}, \frac{M_x}{M} \right).$$

Example 1

Let R be the unit square, $R = \{(x, y) : 0 \leq x \leq 1, 0 \leq y \leq 1\}$. Suppose the density of R is given by the function

$$\rho(x, y) = \frac{1}{y+1}$$

so that R is denser near the x -axis. As a result, we would expect the center of mass to be below the geometric center, $(1/2, 1/2)$. However, since the density does not depend on x , we do expect $\bar{x} = 1/2$.

We have:

$$M = \iint_R \frac{1}{y+1} dA = \int_0^1 \int_0^1 \frac{1}{y+1} dy dx = \int_0^1 \ln(y+1) \Big|_0^1 dx = \int_0^1 \ln 2 dx = \ln 2 = 0.693147\dots$$

$$\begin{aligned} M_x &= \iint_R \frac{y}{y+1} dA = \int_0^1 \int_0^1 \left(1 - \frac{1}{y+1}\right) dy dx = \int_0^1 (y - \ln(y+1)) \Big|_0^1 dx \\ &= \int_0^1 (1 - \ln 2) dx = 1 - \ln 2 = 0.306852819\dots \end{aligned}$$

$$M_y = \iint_R \frac{x}{y+1} dA = \int_0^1 \int_0^1 \frac{x}{y+1} dy dx = \int_0^1 x \ln 2 dx = \frac{1}{2} x^2 \ln 2 \Big|_0^1 = \frac{1}{2} \ln 2 = 0.346573590\dots$$

Thus the center of mass is

$$(\bar{x}, \bar{y}) = \left(\frac{M_y}{M}, \frac{M_x}{M} \right) = \left(\frac{\frac{1}{2} \ln 2}{\ln 2}, \frac{1 - \ln 2}{\ln 2} \right) = \left(\frac{1}{2}, 0.442095\dots \right).$$

Example 2 (Polar)

Let $0 \leq z \leq 1$. Let R be the polar region

$$R = \{(r, \theta) : z \leq r \leq 1, 0 \leq \theta \leq \frac{\pi}{2}\}.$$

Suppose R has constant density ρ . Then:

$$M = \iint_R \rho dA = \rho \iint_R dA = \rho \cdot \text{area of } R = \rho \left(\frac{\pi}{4} - \frac{\pi z^2}{4} \right) = \frac{\pi}{4} \rho (1 - z^2).$$

$$M_x = \iint_R \rho y dA = \rho \int_z^1 \int_0^{\pi/4} r^2 \sin \theta d\theta dr = \rho \int_z^1 -r^2 \cos \theta \Big|_0^{\pi/2} dr = \rho \int_z^1 r^2 dr = \frac{1}{3} \rho (1 - z^3).$$

$$M_y = \iint_R \rho x dA = \rho \int_z^1 \int_0^{\pi/2} r^2 \cos \theta d\theta dr = \rho \int_z^1 r^2 \sin \theta \Big|_0^{\pi/2} dr = \rho \int_z^1 r^2 dr = \frac{1}{3} \rho (1 - z^3).$$

Thus, the center of mass is

$$(\bar{x}, \bar{y}) = \left(\frac{\frac{1}{3}(1 - z^3)}{\frac{\pi}{4}(1 - z^2)}, \frac{\frac{1}{3}(1 - z^3)}{\frac{\pi}{4}(1 - z^2)} \right).$$

An interesting feature of this region is that if z is sufficiently large, the center of mass will be outside the region. This happens when the distance from the center of mass to $(0, 0)$ is less than z . That is, when

$$\sqrt{2} \frac{\frac{1}{3}(1 - z^3)}{\frac{\pi}{4}(1 - z^2)} < z.$$

By factoring, we see that this is equivalent to

$$\frac{\frac{\sqrt{2}}{3}(1 + z + z^2)}{\frac{\pi}{4}(1 + z)} < z.$$

The critical z value is the positive solution to

$$0 = z^2 + z - \frac{\frac{\sqrt{2}}{3}}{\frac{\pi}{4} - \frac{\sqrt{2}}{3}}$$

which is about 0.82337397.... Thus, if $z > 0.82337397...$, the region is very thin, and the center of mass lies outside of the region.

Note:

1. Physical Bodies have Centre of mass, whereas plane bodies have centroid.
2. For physical bodies, density is a function of x and y, whereas for plane bodies, density is assumed to be constant.

Moment of Inertia:

Recall:

$$I_x = \sum m y^2, \text{ MI about X axis}$$

$$I_y = \sum m y x^2, \text{ MI about Y axis}$$

$$I_0 = \sum m (x^2 + y^2) \text{ MI about origin}$$

Using double integrals

$$\text{MI about X axis } I_x = \iint_R y^2 \rho(x, y) dA$$

$$\text{MI about XY axis } I_y = \iint_R x^2 \rho(x, y) dA$$

$$\text{MI about origin } I_0 = \iint_R (x^2 + y^2) \rho(x, y) dA$$